

Medications to Avoid Before and After Surgery

The medications listed below are all medications which may prolong bleeding at the time of your surgery. If you are taking any medications on this list, they should be discontinued 14 days prior to surgery and only Tylenol should be taken for pain. All other medications that you are currently taking must be specifically cleared by your doctor prior to surgery. It is absolutely necessary that all of your current medications be specifically cleared by your doctor and the nursing staff.

4-Way Cold Tabs	5-Aminosalicylic Acid	Acetilsalicylic Acid
Adprin-B products	Alka-Seltzer products	Amigesic
Anacin products	Anexsia w/ Codine	Argesic-SA
Arthra-G	Arthriten products	Arthritis Foundation products
Arthritis Pain Formula	Arthritis Strength BC Powder	Arthropan
ASA	Asacol	Ascriptin products
Aspergum	Asprimox products	Axotal
Azdone	Azulfidine products	B-A-C
Backache Maximum Strength Relief	Bayer products	BC Powder
Bismatrol products	Buffered Aspirin	Bufferin products
Buffetts 11	Buffex	Butal/ASA/Caff
Butalbital Compound	Cama Arthritis Pain Reliever	Carisoprodol Compound
Cheracol	Choline Magnesium Trisalicylate	Choline Salicylate
Cope	Coricidin	Cortisone Medications
Damason-P	Darvon Compound-65	Darvon/ASA
Dolobid	Dristan	Duragesic
Easprin	Ecotrin products	Empirin products
Equagesic	Excedrin products	Fiorgen PF
Fiorinal products	Gelpirin	Genprin
Gensan	Goody's Extra Strength Headache Powders	Halfprin products
Isollyl Improved	Kaodene	Lanorinal
Lortab ASA	Magan	Magnaprin products
Magnesium Salicylate	Magsal	Marnal
Marthritic	Meprobamate	Mesalamine
Methocarbamol	Micrinin	Mobidin
Mobigesic	Momentum	Mono-Gesic
Night-Time Effervescent Cold	Norgesic Products	Norwich products
Olsalazine	Orphengesic products	Oxycodone